Pre-Operative Instruction Checklist

Your Surgery Date is:	
Duncariutions Donairead	Antibiotic: Pain Killer: Norco/Vicoden Other:
Prescriptions Received: One Month Before Surgery	Antibiotic: Pain Killer: Norco/Vicoden Other:
One Worth Before Surgery	Refrain from all nicotine products
3 Weeks Before Surgery	Netrain from all flicotiffe products
	Stop Aspirin or any products that contain aspirin.
	Stop taking Advil, Motrin, Ibuprofen, Aleve, or any medication that contains those
	products.
	*** Tylenol (Acetaminophen)is the only medication acceptable 3 weeks prior to
	your surgery and for one week after surgery.***
	If over the age of 50, EKG is required within last year.
1 Week Before Surgery	
	Do not drink Alcohol for 1 week before and 1 week after your surgery.
1 D B - (C	Stop all herbal remedies unless approved by Dr. Evangelisti
1 Day Before Surgery	Nous pand to stant an authorized at a d
	You need to start your antibiotics today. The suggestion and the start your beauty and the start of the
	 The surgery center will contact you to let you know when you are to arrive to at the surgery center for your procedure. If you are scheduled for Friday, they will
	call you on Thursday. If you are scheduled for Monday, they will call you on
	Friday.
	DO NOT EAT OR DRINK ANYTHING AFTER MIDNIGHT!!!! **No mints, candies, or
	gum, ice chips, or water.
Day of Surgery	
	 You may shower and shampoo. Please do not use any make-up, hair products,
	body/facial moisturizers, deodorant or dark nail polish.
	Wear clothing that either buttons up or zips up.
	If you are an augment or breast reconstructive surgery patient, take your 2 Florest tablets with a six of water an augment to the augment and the surgery patient.
	 Flexeril tablets with a sip of water on your way to the surgery center Do not take your antibiotics today as you will receive antibiotics at the surgical
	center.
	 You may take the following medications the morning of surgery with a small sip of
	water:
Day After Surgery	
	 Resume your antibiotics today. Finishing off the next 9 days.
	 Make sure to take a lot of fluids and eat prior to taking your pain medication.
	Antibiotics have a refill on them. If you have a drain in place longer than 9 days,
	refill antibiotics and take them until 2 days <u>after</u> drain is removed.
After surgery	
	Arnica Montana (6C) will help to resolve bruising. Dissolve 3-5 pellets under
	tongue 3 times/day.
	Narcotic can cause constipation. Please use magnesium citrate to alleviate
	symptoms.
	Please call the office with any questions/concerns prior to upcoming weekend.